

Reply to ‘‘Comment on ‘Phenomenological damping in optical response tensors’’’

A. D. Buckingham and P. Fischer

Department of Chemistry, University of Cambridge, Lensfield Road, Cambridge CB2 1EW, United Kingdom

(Received 19 September 2000; published 16 March 2001)

We show that damping factors must not be incorporated in the perturbation of the ground state by a static electric field. If they are included, as in the theory of Stedman *et al.* [preceding Comment, Phys. Rev. A **63**, 047801 (2001)], then there would be an electric dipole in the y direction induced in a hydrogen atom in the $M_S = +\frac{1}{2}$ state by a static electric field in the x direction. Such a dipole is excluded by symmetry.

DOI: 10.1103/PhysRevA.63.047802

PACS number(s): 42.50.Ct, 03.65.-w, 33.80.-b, 78.20.Jq

We note that Stedman *et al.* (SNAD) [1] now agree that optical susceptibilities in the polarization formalism should have the signs of damping factors as in Bloembergen [2,3], Flytzanis [4], Shen [5], Butcher and Cotter [6], and our paper [7]. An important issue remaining, however, is the question of whether damping is ‘‘inevitable from any perturbation, including a static field’’ [1]. Andrews *et al.* [8] and SNAD assert that it is inevitable. We shall address this important issue by considering the response of a spherical atom to a static electric field and shall show that the assertion is false.

Suppose that damping is necessary in the description of the first-order perturbed ground-state ket of an atom in a

static electric field F_β . Such a ket may then be written in the form

$$|g_F^{\text{damp}}\rangle = |g\rangle + \sum_{j \neq g} \frac{\langle j | \hat{\mu}_\beta | g \rangle}{\hbar \left(\omega_{jg} - \frac{i}{2} \Gamma_{jg} \right)} |j\rangle F_\beta, \quad (1)$$

where $\hat{\mu}$ is a dipole operator, ω_{jg} is the transition angular frequency between the upper level j and the ground state g , and Γ_{jg} is the associated damping factor. The summation is over all excited states $j \neq g$. The expectation value of the α component of the atom’s dipole moment in the field would then be

$$\begin{aligned} \mu_\alpha &= \langle g_F^{\text{damp}} | \hat{\mu}_\alpha | g_F^{\text{damp}} \rangle = \sum_{j \neq g} \left\{ \frac{\langle g | \hat{\mu}_\alpha | j \rangle \langle j | \hat{\mu}_\beta | g \rangle}{\hbar \left(\omega_{jg} - \frac{i}{2} \Gamma_{jg} \right)} + \frac{\langle g | \hat{\mu}_\beta | j \rangle \langle j | \hat{\mu}_\alpha | g \rangle}{\hbar \left(\omega_{jg} + \frac{i}{2} \Gamma_{jg} \right)} \right\} F_\beta \\ &= \sum_{j \neq g} \frac{2\omega_{jg} \text{Re}\{\langle g | \hat{\mu}_\alpha | j \rangle \langle j | \hat{\mu}_\beta | g \rangle\} - \Gamma_{jg} \text{Im}\{\langle g | \hat{\mu}_\alpha | j \rangle \langle j | \hat{\mu}_\beta | g \rangle\}}{\hbar \left(\omega_{jg}^2 + \frac{1}{4} \Gamma_{jg}^2 \right)} F_\beta. \end{aligned} \quad (2)$$

If $|g\rangle$ is degenerate, as for a hydrogen or sodium atom in an $M_S = \pm \frac{1}{2}$ spin state, then $|g\rangle$ is complex, and both the real and imaginary parts of $\langle g | \hat{\mu}_\alpha | j \rangle \langle j | \hat{\mu}_\beta | g \rangle$ are nonzero. The imaginary part changes sign on interchanging α and β . This would imply that a static field F_x induces a dipole in the y direction in a spherical atom in an $M_S = +\frac{1}{2}$ state. However, this is incompatible with quantum theory, for such a dipole must be zero by symmetry (it would be equal and opposite for the $M_S = +\frac{1}{2}$ and $-\frac{1}{2}$ states and therefore change sign under time reversal).

If $|g\rangle$ is nondegenerate, as for a helium atom, then $|g\rangle$ may, without loss of generality, be chosen to be real. The induced dipole would be

$$\mu_x = \sum_{j \neq g} \frac{2\omega_{jg} |\langle g | \hat{\mu}_x | j \rangle|^2}{\hbar^2 \left(\omega_{jg}^2 + \frac{1}{4} \Gamma_{jg}^2 \right)} F_x = \alpha F_x, \quad (3)$$

which is an incorrect equation for the static polarizability α . The correct equation for α is obtained from Eq. (3) by omitting all Γ_{jg} .

We note that SNAD appear to have missed the notes in the epilogue to Bloembergen’s book [2]; the section SNAD refer to has been replaced by the following statement: ‘‘The correct limiting behavior for the case that either the electromagnetic frequency or the material resonant frequency becomes very small, $\omega \rightarrow 0$ or $\omega_{ng} \rightarrow 0$, respectively, requires a more careful treatment of the damping terms, as has been discussed in detail by Van Vleck and Weisskopf [9].’’

SNAD do not refute our argument that inclusion of damping leads to a complex ket $|g_F^{\text{damp}}\rangle$ even for a nondegenerate ground state, such that $|g_F^{\text{damp}}\rangle$ is linearly independent of $|g_F^{\text{damp}}\rangle^*$, which is incompatible with nondegeneracy.

We conclude that damping factors must not be incorporated in the perturbation of the ground state by a static field.

- [1] G. E. Stedman, S. Naguleswaran, D. L. Andrews, and L. C. Dávila Romero, preceding Comment, Phys. Rev. A **63**, 047801 (2000).
- [2] N. Bloembergen, *Nonlinear Optics*, 4th ed. (World Scientific, Singapore, 1996).
- [3] N. Bloembergen, H. Lotem, and R. T. Lynch, Indian J. Pure Appl. Phys. **16**, 151 (1978).
- [4] C. Flytzanis, in *Quantum Electronics*, edited by H. Rabin and C. L. Tang (Academic, New York, 1975), Vol. 1, Pt. A.
- [5] Y. R. Shen, *The Principles of Nonlinear Optics* (John Wiley & Sons, New York, 1984).
- [6] P. N. Butcher and D. Cotter, *The Elements of Nonlinear Optics, Cambridge Studies in Modern Optics*, reprint ed. (Cambridge University Press, Cambridge, 1993).
- [7] A. D. Buckingham and P. Fischer, Phys. Rev. A **61**, 035801 (2000).
- [8] D. L. Andrews, S. Naguleswaran, and G. E. Stedman, Phys. Rev. A **57**, 4925 (1998).
- [9] J. H. Van Vleck and V. F. Weisskopf, Rev. Mod. Phys. **17**, 227 (1945).